

Procedures / Interventions	Med Doses	EMT	AEMT	PM
<p>• Perform initial assessment and treatments as described in IPAP.</p> <p>A patient may be identified as being at risk for Adrenal Insufficiency or Adrenal Crisis by presence of one of the following:</p> <ul style="list-style-type: none"> ➤ Medical alert bracelet/identification ➤ Patient medical records ➤ History reported by patient or family <p>Patients with adrenal insufficiency who present with any of the following signs/symptoms may need “stress dose” steroids if available.</p> <ul style="list-style-type: none"> ➤ Severe injury of any type ➤ Acute blood loss ➤ Fluid and Electrolyte loss ➤ Infections ➤ Severe Nausea/Vomiting/Diarrhea ➤ Abdominal pain ➤ Altered mental status <p>Treatment:</p> <ul style="list-style-type: none"> ☎ If patient or family member has physician-prescribed “stress dose” steroid medication available, contact medical control for orders to administer steroid. ➤ Obtain vascular access IV/IO if appropriate ➤ Aggressive fluid resuscitation may be necessary during transport. ➤ Assess blood glucose if indicated and treat according to the <u>Hypoglycemia and Hyperglycemia Protocol</u> 				<p>•</p> <p>•</p> <p>•</p> <p>•</p>